

Literaturverzeichnis zu:

Detlef Kampf:

Optimale Kupferversorgung:

Detlef Kampf:

Optimal copper supply



In **07-08 /2012, S. 20 ff**

1. **Allee et al. (2011)**: Effects of Feeding Tribasic Copper Chloride or Copper Sulfate on Growth and Efficiency of Nursery Pigs. Proc. Trace mineral conference, Munich.
2. **Banks et al. (2004)**: The effects of copper on the efficacy of phytase, growth, and phosphorous retention in broiler chicks. Poul. Sci., 83, 1335-1341.
3. **EC Regulation 1334/2003**: Conditions for authorisation of a number of additives in feedingstuffs belonging to the group of trace elements. 25 July 2003.
4. **Helle and Kampf (2008)**: Analytical possibilities to differentiate Trace Element-Amino Acid-Chelates. In: Schlegel, P., Durosoy, S., Jongbloed, A.W. (eds.) Trace elements in animal production systems. Wageningen Academic Publishers, Wageningen, 247-249.
5. **Kincaid et al. (1997)**: Zinc Oxide and Amino Acids as Sources of Dietary Zinc for Calves: Effects on Uptake and Immunity. J Dairy Sci 80:1381–1388
6. **Liu et al. (2005)**: Layer performance and phytase retention as influenced by copper sulfate pentahydrate and tribasic copper chloride. J. Appl. Poul. Res., 14(3), 499-505.
7. **Lu et al. (2010)**: Effect of Dietary Supplementation with Copper Sulfate or Tribasic Copper Chloride on the Growth Performance, Liver Copper Concentrations of Broilers Fed in Floor Pens, and Stabilities of Vitamin E and Phytase in Feeds. Biol. Trace Elem. Res., 138 (1-3), 181-189.
8. **Luo et al. (2005)**: Effects of dietary supplementation with copper sulfate or tribasic copper chloride on broiler performance, relative copper bioavailability, and oxidation stability of vitamin E in feed. Poul. Sci., 84, 888-893.
9. **Miles et al. (1998)**: The Effect of dietary supplementation with copper sulfate or tribasic copper chloride on broiler performance, relative copper bioavailability, and dietary prooxidant activity. Poul. Sci., 77, 416-425.
10. **Nockels et al. (1993)**: Stress Induction Affects Copper and Zinc Balance in Calves Fed Organic and Inorganic Copper and Zinc Sources. J. Anim. Sci., 71, 2539-2545.
11. **Pang and Applegate (2006)**: Effects of Copper Source and Concentration on in Vitro Phytate Phosphorus Hydrolysis by Phytase. J. Agric. Food Chem., 54, 1792-1796.

12. **Spears et al. (2004)**: Bioavailability of copper from tribasic copper chloride and copper sulfate in growing cattle. *Anim. Feed Sci. Technol.*, 116, 1–13.
13. **Wedekind et al. (1992)**: Methodology for Assessing Zinc Bioavailability: Efficacy Estimates for Zinc-Methionine, Zinc Sulfate and Zinc Oxide. *J. Anim. Sci.*, 70, 178-187.